

ERRATUM

IN SEARCH OF A DIABETIC CARDIOPATHY — A STUDY OF LEFT VENTRICULAR FUNCTION IN 25 ASYMPTOMATIC DIABETICS USING M MODE ECHOCARDIOGRAPHY — A PILOT STUDY

by S Balasingam, PK Mah, C H Lee
SMJ 1988; 29: 246-251

Table 1 was omitted and is reproduced as follows:

TABLE 1: ECHOCARDIOGRAPHIC DATA IN 25 ASYMPTOMATIC DIABETICS IN RELATION TO DURATION, TYPE OF DIABETES AND GLYCAEMIC CONTROL

	S/NO	FS %	EF %	LA/AO	DURATION IN YEARS	HbA1c	TYPE
1	11	8	23	1.50	30	13.3	IDDM
2	17	15	40	0.89	1	8.3	NIDDM
3	90	16	41	1.00	8	13.9	IDDM
4	20	18	38	0.96	14		NIDDM
5	13	18	46	1.50	22	16.5	IDDM
6	01	19	47	1.20	8		IDDM
7	04	19	47	1.00	10	11.5	IDDM
8	12	19	47	1.34	8	7.3	IDDM
9	21	20	43	1.00	3		IDDM
10	16	21	50	1.27	14	16.6	IDDM
11	10	22	53	0.96	10	13.3	IDDM
12	08	22	53	1.21	4	13.6	NIDDM
13	24	22	53	0.81	3	10.4	NIDDM
14	14	22	54	0.87	10	14.8	IDDM
15	15	23	40	0.75	2	17.6	IDDM
16	06	23	55	0.83	1		IDDM
17	03	27	62	1.30	10	7.7	NIDDM
18	18	28	64	0.93	1		NIDDM
19	02	29	64	1.03	6	10.7	NIDDM
20	23	29	64	0.96	10	12.3	NIDDM
21	05	31	67	0.83	3		IDDM
22	07	33	70	1.00	10	12.1	NIDDM
23	25	37	75	1.17	2	13.2	NIDDM
24	19	39	72	0.50	4	7.7	NIDDM
25	22	42	74	0.95	12		NIDDM

The data set is sorted according to FS, from lowest to highest.

Note that 16/25 had FS < 25%, and 9/25 had FS > 25%.

Of those with FS 25, all had NIDDM except one.

NIDDM : non insulin dependent diabetes

IDDM : insulin dependent diabetes

HbA1c : glycosylated haemoglobin

FS : fractional shortening

LA : left atrium

AO : aortic root

S/NO : serial number of patient

EF : ejection fraction

Local Normal Values

FS : 26% — 39%

LA/AOR: 0.86 — 1.4

HbA1c : 6.4 — 9.6%

EF : 60 — 77%