

MALIGNANT RECTAL PERFORATION?

Dear Sir,

I read the article, "Traumatic rectal perforation presenting as necrotising fasciitis of the lower limb" with great interest. The authors have described the clinical presentation, diagnosis and management explicitly. However, I have some queries.

Firstly, the distance between the site of the perforation and the previous anastomotic site was not mentioned. It may possibly be a rupture of the anastomotic line due to a recurrence. Secondly, it is also not known whether the biopsy was taken from the perforation site or not. Without a biopsy, the possibility of a recurrence cannot be excluded. Various studies have reported different rates of recurrence in carcinoma of the rectum.⁽¹⁻³⁾ Galandink et al have mentioned a recurrence rate as high as 52% within a time interval of one month and seven-and-a-half years.⁽²⁾ Lastly, was the loop colostomy ever closed? The authors have mentioned a two-year follow-up without providing more information about the fate of the colostomy.

Yours sincerely,

Nikhil Gupta

Niladhar Shankarrao Hadke

Department of Surgery
Maulana Azad Medical College
Bahadur Shah Zafar Marg
Delhi 110002
India
Email: nikhil_ms26@yahoo.co.in

REFERENCES

1. Fu WP, Quah HM, Eu KW. Traumatic rectal perforation presenting as necrotising fasciitis of the lower limb. *Singapore Med J* 2009; 50: e270-e273.
2. HIMAL HS. Anastomotic recurrence of carcinoma of the colon and rectum. The value of endoscopy and serum CEA levels. *Am J Surg* 1991; 57:334-7.
3. Galandink S, Wieand HS, Moertel CG, et al. Patterns of recurrence after curative resection of carcinoma of the colon and rectum. *Surg Gynecol Obstet* 1992; 174:27-32.
4. Polglase AL, Grodski SF, Tremayne AB, Chee JB, Bhathal PS. Local recurrence following surgical treatment for carcinoma of the lower rectum. *ANZ J Surg* 2004; 74:745- 50.