

Improving the quality of CPR in the community

Ong E H M

ABSTRACT

Bystander cardiopulmonary resuscitation (CPR) is important for survival from out-of-hospital cardiac arrest (OHCA). However, recent research indicates that the quality of CPR is an important and often overlooked factor affecting survival. Individual factors, training, awareness, technique and rescuer fatigue may influence the quality of CPR. Quality components of CPR include rate, ratio, depth and ventilation-compression ratio. The new 2010 CPR guidelines advocate a ratio of compressions to ventilations of 30:2, with a rate of at least 100 compressions per minute. Depth of compression should be at least 5 cm. Rescuers should allow complete recoil of the chest. Locally, limited information is available regarding the quality of CPR being performed for OHCA. Strategies to improve the quality of CPR include research, training, education as well as incorporating appropriate technologies that measure and feedback the quality of CPR. These technologies are at the heart of recent advances, as they now make it feasible to provide routine feedback to rescuers providing CPR, through the integration of feedback devices into training equipment, defibrillators and standalone CPR assist devices.

Keywords: bystander, cardiac arrest, cardiopulmonary resuscitation, community, quality

Singapore Med J 2011;52(8):586-591

INTRODUCTION

Of the approximately 16,000 deaths that occur in Singapore every year, about 23% are from a cardiac cause,⁽¹⁾ some of which will occur suddenly and outside of a hospital. A study conducted from October 2001 to September 2004 found that a total of 2,428 out-of-hospital cardiac arrests (OHCA) occurred in Singapore.⁽²⁾ The mechanism of death is usually a fatal arrhythmia, most often ventricular tachycardia or fibrillation.⁽³⁾ Many of these patients would be clinically dead long before they are transported to a hospital. In the chain of survival concept,^(4,5) the provision

of early bystander cardiopulmonary resuscitation (CPR) in OHCA is important to improve survival in sudden cardiac arrest. There is currently good evidence that indicates the importance of CPR combined with early defibrillation (< 4 mins).⁽⁶⁻⁹⁾ Survival rates for pre-hospital cardiac arrest vary in published reports, from 2% to over 20% in various cities and countries.⁽¹⁰⁾

Singapore's Emergency Medical Services (EMS) system is run by the Singapore Civil Defence Force (SCDF), which currently operates 36 ambulances based in 14 fire stations and more than ten fire posts in a single-tier system. Private ambulance operators do not generally convey emergency cases. Singapore EMS is activated by a universal, centralised and enhanced '995' dispatching system run by the SCDF, which utilises a computer-aided dispatch system, medical dispatch protocols, Global Positioning Satellite automatic vehicle locating systems and road traffic monitoring systems. Since 1996, ambulances in Singapore have been manned by specifically trained paramedics. They are able to provide Basic Life Support (BLS) and defibrillation with automated external defibrillators (AED), intravenous adrenaline and saline infusions. They also perform laryngeal mask airway insertion in cardiac arrest.

The Cardiac Arrest and Resuscitation Epidemiology (CARE) study group includes representatives from the six major public hospitals in Singapore, the SCDF, Health Sciences Authority and the Clinical Trials and Epidemiology Research Unit of the Ministry of Health, Singapore. The CARE phase I study, which described the OHCA epidemiology in Singapore,⁽¹¹⁾ found survival from OHCA in Singapore to be 2.0%. The mean and standard deviation of EMS response time was found to be 10.2 ± 4.3 minutes and that of time from call to defibrillation was 16.7 ± 7.2 minutes. 35.3% of arrests were not witnessed, 54.6% were bystander witnessed and 10.0% were witnessed by EMS personnel. Bystander CPR was present for 20.6% of all cases, and a further 10.0% received immediate CPR from EMS personnel who witnessed the arrest.⁽¹¹⁾

QUALITY OF CPR

Previous studies have demonstrated the importance of bystander CPR to survival from OHCA.^(7,12) However, recent

Department of Emergency Medicine, Singapore General Hospital, Outram Road, Singapore 169608

Ong EHM, FRCSE, MPH, FAMS
Consultant and Senior Medical Scientist

Correspondence to:
Dr Ong Eng Hock
Marcus
Tel: (65) 6321 3590
Fax: (65) 6321 4873
Email: marcus.ong.e.h@sgh.com.sg

Table I. International Liaison Committee on Resuscitation recommendations for quality of adult CPR.

	ILCOR 2005 ⁽³¹⁾	ILCOR 2010 ⁽³²⁾
Ventilation rate	8–10/min	No change
Inspiratory time	< 1 sec	No change
Tidal volume	500–600 ml	No change
Compression rate	> 100/min	No change
Compression depth	4–5 cm	> 5 cm
Chest recoil	50% duty cycle complete recoil (compression/ release)	No change
Compression: ventilation ratio	30:2	No change

CPR: cardiopulmonary resuscitation

research has indicated that the quality of CPR is an important but often overlooked factor affecting survival.^(13,14) The current International Liaison Committee on Resuscitation (ILCOR) Advanced Cardiac Life Support Guidelines (2005)^(15,16) suggest that the quality of CPR is as important as the presence or absence of bystander CPR.

Several factors may influence the quality of CPR, including individual factors, training, awareness, technique as well as rescuer fatigue. Quality components of CPR include the rate, ratio, depth as well as ventilation-compression ratio. There is good evidence to show that excessive ventilation rates during CPR result in significantly increased intrathoracic pressures and markedly decreased coronary perfusion pressures.⁽¹⁷⁻¹⁹⁾ It is suggested that hyperventilation frequently occurs during CPR, even when performed professional rescuers,⁽¹⁸⁾ and survival rates decrease as a result. The 2005 CPR guidelines^(16,20) recommended an inspiratory time of one second, at a rate of 8–10 ventilations per minute. In a patient with an established advanced airway (e.g. endotracheal tube), it is reasonable to deliver ventilations without pausing chest compressions.

Another important factor is the rate of chest compressions and the problem of prolonged interruption of chest compressions during CPR, resulting in less cycle time on chest compression and lower coronary perfusion pressures.⁽²¹⁻²⁴⁾ Studies have shown that the rates of chest compression during CPR are often far less than those recommended.^(14,25,26) One study has shown improved haemodynamics in humans with high-frequency CPR (120/min) compared with standard CPR.⁽²⁷⁾ The depth of compression achieved has also been shown in clinical studies to be often much less than that recommended by CPR^(25,28) guidelines. In animal models, the depth of compression was related to better outcomes.⁽²⁹⁾ Also, incomplete chest recoil has been found to be associated with increased intrathoracic

pressure, resulting in decreased coronary and cerebral perfusion.⁽³⁰⁾

The local 2006 CPR guidelines⁽²⁰⁾ advocate a 30:2 ratio of compressions to ventilations, with a rate of 100 compressions per minute. Depth of compression should be at least 4–5 cm. Rescuers should also allow complete recoil of the chest, with a ratio of compression to release of 50%. Table I shows a comparison between the 2005 and 2010 ILCOR guidelines for quality of CPR.^(31,32)

Researchers have been using the following parameters to measure the quality of CPR: rate of chest compressions; rate of ventilations; and CPR flow fraction, usually for the first five minutes of resuscitation. The CPR flow fraction and its counterpart, the ‘no flow ratio’ (NFR), reflect the ratio of compressions to pauses (‘no flow’) in the CPR cycle.⁽³³⁾ Compression is usually defined as the fraction of time with sub-zero position of the sternum, while ‘no flow’ is defined as all pauses between compressions longer than 1.5 seconds. The sum of such intervals is then divided by the segment length (e.g. five minutes), from which the NFR can be derived. These research parameters are usually obtained automatically using various software programmes bundled with the latest defibrillators. Locally, there is currently limited information available regarding the quality of CPR being performed for OHCA in Singapore. In the CARE phase I study, bystander CPR was present for 20.6% (103 out of 500) of all cases; however, only 67.0% of these (69 out of 103) were reported to have received CPR with both compression and ventilation. Another 26.2% (27 out of 103) received chest compression only.⁽¹¹⁾

Regarding the Emergency Department, a study on the quality of CPR found significant interruptions to CPR with both manual and mechanical CPR.⁽³⁴⁾ The ‘no flow time’ (NFT) (defined as the sum of all pauses between compressions longer than 1.5 seconds) during the first five minutes of resuscitation was manual CPR 85 seconds vs. mechanical CPR 104 seconds (difference = 19 seconds; 95% confidence interval [CI] –18 to 56). The NFR, defined as NFT divided by segment length, was manual 0.28 vs. LDB 0.40 (difference = –0.12; 95% CI –0.22 to –0.02).

IMPROVING THE QUALITY OF CPR

Several strategies have been suggested in order to improve the quality of CPR in our community. These include research, training, education and incorporating appropriate technologies that measure and feedback the quality of CPR. These technologies are at the heart of recent advances, as they now make it feasible to provide routine feedback to rescuers providing CPR through




Fig. 1 Photograph shows defibrillator pads incorporating an accelerometer for QCPR feedback (Zoll E series defibrillation pads).



Fig. 2 Defibrillator screen display incorporating QCPR feedback indicators (Zoll E series). When complete filling of the CPR compression indicator has not been achieved due to diminished compression rate or depth, the E Series will display the letter R for rate and/or the letter D for depth to assist the rescuer in determining whether chest compression rate or depth should be increased. When appropriate rate and depth have been achieved, these letters will disappear from the display field. A voice prompt option is also available.




Fig. 3 QCPR feedback showing the number of compressions and ventilations per minute over a period (Medtronic-Physiocontrol Codestat Suite).

integration of feedback devices into training equipment, defibrillators as well as standalone CPR assist devices.

CPR feedback devices have long been incorporated into training manikins used for CPR instruction. These are able to provide feedback to trainees regarding the rate and depth of chest compressions, as well as the rate and volume of ventilations. Training using a manikin equipped with a computer-based voice advisory feedback system has been shown to improve retention of CPR skills.⁽³⁵⁾ The use of a video recording of a trainee’s performance is now an additional tool available for instruction. It cannot be over-emphasised that CPR training centres should keep up-to-date with the latest CPR guidelines and ensure good quality CPR feedback during instruction, so that proper techniques can be learnt and retained.⁽³⁶⁾ This also applies to healthcare providers as well as public teaching of CPR. This attention to quality also highlights the need for accreditation of CPR training centres and regular audits

in order to maintain high standards. It is also important to ensure trainees are kept updated on the latest CPR guidelines through recertification.

More importantly, devices to facilitate real-time feedback to CPR providers are now available. The latest generation of defibrillators utilise accelerometers embedded within chest defibrillation pads to measure the depth and rate of compression, as well as variations in chest impedance that can reflect chest wall movements.^(14,25) Fig. 1 shows an example of a device that incorporates an accelerometer, which is located on the sternum, beneath the heel of the hand, while manual compressions are going on. These devices are able to give verbal as well as visual prompts to rescuers performing CPR, and cue the rescuer to speed up, slow down or increase the depth of compressions or ventilations.⁽³⁷⁾ An example of these visual cues is shown in Fig. 2. Such devices have been shown to improve the quality of CPR for OHCA⁽³³⁾ as well as in-hospital cardiac




Fig. 4 (a & b) Photographs show a pocket Q CPR feedback device for use with manual CPR (CPREZY).

arrest (IHCA).⁽³⁷⁾ Provision should be made for the gradual upgrading of existing defibrillators to include quality of CPR (QCPR) technology in the hospital and in out-of-hospital settings. Fig. 3 shows an example of the QCPR summary data available from such devices. Small, pocket-sized, standalone devices are also now commercially available to provide rescuers with real-time CPR feedback (Fig. 4). Education of the public will help increase the usage of such devices among lay rescuers.

In the hospital setting, collection of CPR data should become a routine part of hospital quality assurance processes. Adoption of defibrillators using QCPR software will enable routine monitoring, evaluation and feedback to CPR providers from IHCA. For example, the National Registry of Cardiopulmonary Resuscitation (NRCPR) is an American Heart Association (AHA)-sponsored, prospective, multi-site, observational study of in-hospital resuscitation.⁽³⁸⁾ QCPR data is one of the items routinely collected by study sites.

Good quality research on QCPR in our community is much needed locally in order to identify areas that require intervention. Innovative strategies are needed both to increase the number of trained CPR providers in Singapore as well as the quality of instruction and CPR performed. For example, it has been suggested that widespread video-based CPR instruction may be used as a simple tool to teach CPR,⁽³⁹⁾ so as to reach a larger group of the population. Others have looked at identifying targeted, high-risk groups for CPR instruction.⁽⁴⁰⁾ Another area for implementing QCPR is with dispatcher-assisted CPR.⁽⁴¹⁾ This occurs when the EMS dispatcher is able to give instructions on how to perform CPR to a willing rescuer over the telephone. QCPR technology can now be integrated with dispatcher CPR protocols^(42,43) to cue the rescuer as to how many compressions and ventilations to give per minute, as the rescuer counts out with the dispatcher over the telephone. A compression and ventilation rate counter is integrated with

the dispatcher's tools on his computer screen. There have been proposals to upgrade Singapore's dispatch centre with this technology.

A more controversial area is the use of chest compression-only CPR (CC-CPR) without ventilations as an alternative to standard CPR for bystanders.⁽⁴⁴⁾ Proponents of CC-CPR argue that it overcomes bystander reluctance to do mouth-to-mouth ventilations,⁽⁴⁵⁻⁵³⁾ is simpler to teach⁽⁵⁴⁻⁵⁶⁾ and may result in fewer interruptions to chest compressions.⁽²¹⁻²⁴⁾ However, there have been controversies regarding the relative effectiveness of CC-CPR.⁽⁵⁷⁻⁵⁹⁾ Current ILCOR guidelines state that CC-CPR should be encouraged only if the rescuers are unwilling to do airway or breathing manoeuvres, or are untrained in CPR and are thus uncertain about how to perform CPR. These guidelines also recognise the need for more research on the efficacy of CC-CPR.

Evidence from swine models of cardiac arrest suggests that ventilation may not be essential in the initial 12 minutes of resuscitation^(23,60) and that oxygen tensions can be maintained above 100 mmHg with only CC-CPR.⁽⁶¹⁾ However, when the animal model was simulated with an obstructed airway, as might often be the case in real-life cardiac arrest, arterial blood was completely desaturated within two minutes.⁽⁶²⁾ Also, in a human model of cardiac arrest, passive ventilation during CC-CPR was limited in its ability to maintain adequate gas exchange.⁽⁵⁸⁾

Finally, with the development of portable, field-deployable, mechanical CPR devices,⁽⁶³⁻⁶⁵⁾ there is potential to overcome the problems of rescuer fatigue and individual variations in the quality of CPR. More research and development are needed in order to produce cost-effective mechanical CPR devices that can be widely mass produced.

IMPLICATIONS FOR OUR COMMUNITY

We now know that encouraging community bystander CPR by itself is insufficient. More emphasis on QCPR

in order to improve survival rates in our community is required. This translates into an important message for community CPR education.⁽⁶⁶⁾ Emphasis on QCPR should be incorporated into community CPR training, and also into the training and mindset of CPR instructors. These instructors would be the medium through which good quality practices can be passed on to the providers who would be the eventual lifesavers in the community.

Meanwhile, emphasis should also be placed on improving our community EMS response.⁽⁶⁷⁾ This includes continuous quality improvement and research on our EMS dispatch and ambulance services. Efforts should also be made to upgrade and incorporate QCPR technology into defibrillators and other devices used by rescuers. A national registry should be established to monitor the QCPR during both OHCA and IHCA. QCPR data should be collected and used as part of hospital routine quality assurance processes.

The science of resuscitation is constantly evolving as new discoveries emerge. It is inevitable that CPR guidelines will continue to change and be regularly updated and revised. This highlights the importance of having a National Resuscitation Council (NRC) to constantly evaluate the current scientific evidence and adapt the latest knowledge for local use. It also highlights the need for currency and recertification in CPR training. The NRC also has a role in community-wide dissemination of the latest CPR guidelines.

REFERENCES

1. Health facts Singapore. In: Ministry of Health Singapore. [online] Available at: www.moh.gov.sg/mohcorp/statistics.aspx?id=5526. Accessed August 1, 2011.
2. Ong ME, Tan EH, Yan X, et al. An observational study describing the geographic-time distribution of cardiac arrests in Singapore: what is the utility of geographic information systems for planning public access defibrillation? (PADS Phase I). *Resuscitation* 2008; 76:388-96.
3. Bayes de Luna A, Coumel P, Leclercq JF. Ambulatory sudden cardiac death: mechanisms of production of fatal arrhythmia on the basis of data from 157 cases. *Am Heart J* 1989; 117:151-9.
4. Cummins RO, Ornato JP, Thies WH, Pepe PE. Improving survival from sudden cardiac arrest: the "chain of survival" concept. *Circulation* 1991; 83:1832-47.
5. Cummins RO. The "chain of survival" concept: how it can save lives. *Heart Dis Stroke* 1992; 1:43-5.
6. Valenzuela TD, Roe DJ, Nichol G, et al. Outcomes of rapid defibrillation by security officers after cardiac arrest in casinos. *N Engl J Med* 2000; 343:1206-9.
7. Valenzuela TD, Roe DJ, Cretin S, Spaite DW, Larsen MP. Estimating effectiveness of cardiac arrest interventions: a logistic regression survival model. *Circulation* 1997; 96:3308-13.
8. White RD, Asplin BR, Bugliosi TF, Hankins DG. High discharge survival rate after out-of-hospital ventricular fibrillation with rapid defibrillation by police and paramedics. *Ann Emerg Med* 1996; 28:480-5.
9. Eisenberg MS, Copass MK, Hallstrom AP, et al. Treatment of out-of-hospital cardiac arrests with rapid defibrillation by emergency medical technicians. *N Engl J Med* 1980; 302:1379-83.
10. Eisenberg M, Horwood B, Cummins R, Reynolds-Haertle R, Hearne T. Cardiac arrest and resuscitation: a tale of 29 cities. *Ann Emerg Med* 1990; 19:179-86.
11. Ong EHM, Chan YH, Anantharaman V, et al. Cardiac Arrest and resuscitation epidemiology in Singapore (CARE I study). *Prehosp Emerg Care* 2003; 7:427-33.
12. Stiell IG, Wells GA, DeMaio VJ, et al. Modifiable factors associated with improved cardiac arrest survival in a multicenter basic life support/defibrillation system: OPALS Study Phase I results. Ontario Prehospital Advanced Life Support. *Ann Emerg Med* 1999; 33:44-50.
13. Olasveengen TM, Wik L, Steen PA. Quality of cardiopulmonary resuscitation before and during transport in out-of-hospital cardiac arrest. *Resuscitation* 2008; 76:185-90.
14. Wik L, Kramer-Johansen J, Myklebust H, et al. Quality of cardiopulmonary resuscitation during out-of-hospital cardiac arrest. *JAMA* 2005; 293:299-304.
15. ECC Committee, Subcommittees and Task Forces of the American Heart Association: 2005 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 2005; 112 suppl 24:IV1-203.
16. International Liaison Committee on Resuscitation: Part 4: Advanced Life Support. *Resuscitation* 2005; 67:213-47.
17. Aufderheide TP, Lurie KG. Death by hyperventilation: a common and life-threatening problem during cardiopulmonary resuscitation. *Crit Care Med* 2004; 32 Suppl 9:S345-51.
18. Aufderheide TP, Sigurdsson G, Pirralo RG, et al. Hyperventilation-induced hypotension during cardiopulmonary resuscitation. *Circulation* 2004; 109:1960-5.
19. Pepe PE, Raedler C, Lurie KG, Wigginton JG. Emergency ventilatory management in hemorrhagic states: elemental or detrimental? *J Trauma*. 2003; 54:1048-55; discussion 1055-7.
20. Teo WS, Anantharaman V, Lim SH. Update on resuscitation 2006. *Singapore Med J* 2007; 48:100-5.
21. Hostler D, Guimond G, Callaway C. A comparison of CPR delivery with various compression-to-ventilation ratios during two-rescuer CPR. *Resuscitation* 2005; 65:325-8.
22. Kern KB, Hilwig RW, Berg RA, Sanders AB, Ewy GA. Importance of continuous chest compressions during cardiopulmonary resuscitation: improved outcome during a simulated single lay-rescuer scenario. *Circulation* 2002; 105:645-9.
23. Berg RA, Kern KB, Sanders AB, et al. Bystander cardiopulmonary resuscitation. Is ventilation necessary? *Circulation* 1993; 88 (4 pt 1):1907-15.
24. Yu T, Weil MH, Tang W, et al. Adverse outcomes of interrupted precordial compression during automated defibrillation. *Circulation* 2002; 106:368-72.
25. Abella BS, Alvarado JP, Myklebust H, et al. Quality of cardiopulmonary resuscitation during in-hospital cardiac arrest. *JAMA* 2005; 293:305-10.
26. Abella BS, Sandbo N, Vassilatos P, et al. Chest compression rates during cardiopulmonary resuscitation are suboptimal: a prospective study during in-hospital cardiac arrest. *Circulation* 2005; 111:428-34.
27. Swenson RD, Weaver WD, Niskanen RA, Martin J, Dahlberg S. Hemodynamics in humans during conventional and experimental methods of cardiopulmonary resuscitation. *Circulation* 1988; 78:630-9.
28. Wik L, Steen PA, Bircher NG. Quality of bystander cardiopulmonary resuscitation influences outcome after prehospital cardiac arrest. *Resuscitation* 1994; 28:195-203.
29. Babbs CF, Voorhees WD, Fitzgerald KR, Holmes HR, Geddes LA. Relationship of blood pressure and flow during CPR to chest

- compression amplitude: evidence for an effective compression threshold. *Ann Emerg Med* 1983; 12:527-32.
30. Yannopoulos D, McKnite S, Aufderheide TP, et al. Effects of incomplete chest wall decompression during cardiopulmonary resuscitation on coronary and cerebral perfusion pressures in a porcine model of cardiac arrest. *Resuscitation* 2005; 64:363-72.
 31. International Liason Committee on Resuscitation: Part 2: Adult Basic Life Support. *Resuscitation* 2005; 65:187-201
 32. Sayre MR, Koster RW, Botha M, et al. Part 5: Adult basic life support: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation* 2010; 122 Suppl 2: S298-324.
 33. Kramer-Johansen J, Myklebust H, Wik L, et al. Quality of out-of-hospital cardiopulmonary resuscitation with real time automated feedback: a prospective interventional study. *Resuscitation* 2006; 71:283-92.
 34. Ong ME, Annathurai A, Shahidah A, et al. Cardiopulmonary resuscitation interruptions with use of a load-distributing band device during emergency department cardiac arrest. *Ann Emerg Med* 2010; 56:233-41.
 35. Wik L, Myklebust H, Auestad BH, Steen PA. Twelve-month retention of CPR skills with automatic correcting verbal feedback. *Resuscitation* 2005; 66:27-30.
 36. Pamell MM, Larsen PD. Poor quality teaching in lay person CPR courses. *Resuscitation* 2007; 73:271-8.
 37. Abella BS, Edelson DP, Kim S, et al. CPR quality improvement during in-hospital cardiac arrest using a real-time audiovisual feedback system. *Resuscitation* 2007; 73:54-61.
 38. Peberdy MA, Kaye W, Ornato JP, et al. Cardiopulmonary resuscitation of adults in the hospital: a report of 14720 cardiac arrests from the National Registry of Cardiopulmonary Resuscitation. *Resuscitation* 2003; 58:297-308.
 39. Todd KH, Heron SL, Thompson M, et al. Simple CPR: A randomized, controlled trial of video self-instructional cardiopulmonary resuscitation training in an African American church congregation. *Ann Emerg Med* 1999; 34:730-7.
 40. Richman PB, Bobrow BJ, Clark L, Noelck N, Sanders AB. Ability of citizens in a senior living community to perform lifesaving cardiac skills and appropriately utilize AEDs. *J Emerg Med* 2007; 33:395-9.
 41. Hallstrom AP, Cobb LA, Johnson E, Copass MK. Dispatcher assisted CPR: implementation and potential benefit. A 12-year study. *Resuscitation* 2003; 57:123-9.
 42. Deakin CD, Cheung S, Petley GW, Clewlow F. Assessment of the quality of cardiopulmonary resuscitation following modification of a standard telephone-directed protocol. *Resuscitation* 2007; 72:436-43.
 43. Dias JA, Brown TB, Saini D, et al. Simplified dispatch-assisted CPR instructions outperform standard protocol. *Resuscitation* 2007; 72:108-14.
 44. SOS-KANTO study group. Cardiopulmonary resuscitation by bystanders with chest compression only (SOS-KANTO): an observational study. *Lancet* 2007; 369:920-26.
 45. Braslow A, Brennan RT. Layperson CPR: a response to "A reappraisal of mouth-to-mouth ventilation during bystander-initiated cardiopulmonary resuscitation". *Prehosp Disaster Med* 1999; 14:113-4.
 46. Brenner B, Stark B, Kauffman J. The reluctance of house staff to perform mouth-to-mouth resuscitation in the inpatient setting: what are the considerations? *Resuscitation* 1994; 28:185-93.
 47. Brenner BE, Van DC, Lazar EJ, Camargo C. Determinants of physician reluctance to perform mouth-to-mouth resuscitation. *J Clin Epidemiol* 2000; 53:1054-61.
 48. Caves ND, Irwin MG. Attitudes to basic life support among medical students following the 2003 SARS outbreak in Hong Kong. *Resuscitation* 2006; 68:93-100.
 49. Jelinek GA, Gennat H, Celenza T, et al. Community attitudes towards performing cardiopulmonary resuscitation in Western Australia. *Resuscitation* 2001; 51:239-46.
 50. Locke CJ, Berg RA, Sanders AB, et al. Bystander cardiopulmonary resuscitation. Concerns about mouth-to-mouth contact. *Arch Intern Med* 1995; 155:938-43.
 51. Ornato JP, Hallagan LF, McMahan SB, Peeples EH, Rostafinski AG. Attitudes of BCLS instructors about mouth-to-mouth resuscitation during the AIDS epidemic. *Ann Emerg Med* 1990; 19:151-6.
 52. Shibata K, Taniguchi T, Yoshida M, Yamamoto K. Obstacles to bystander cardiopulmonary resuscitation in Japan. *Resuscitation* 2000; 44:187-93.
 53. Taniguchi T, Omi W, Inaba H. Attitudes toward the performance of bystander cardiopulmonary resuscitation in Japan. *Resuscitation* 2007; 75:82-7.
 54. Assar D, Chamberlain D, Colquhoun M, et al. Randomised controlled trials of staged teaching for basic life support. 1. Skill acquisition at bronze stage. *Resuscitation* 2000; 45:7-15.
 55. Lester CA, Donnelly PD, Assar D. Lay CPR trainees: retraining, confidence and willingness to attempt resuscitation 4 years after training. *Resuscitation* 2000; 45:77-82.
 56. Chamberlain D, Smith A, Colquhoun M, et al. Randomised controlled trials of staged teaching for basic life support: 2. Comparison of CPR performance and skill retention using either staged instruction or conventional training. *Resuscitation* 2001; 50:27-37.
 57. Waalewijn RA, Tijssen JG, Koster RW. Bystander initiated actions in out-of-hospital cardiopulmonary resuscitation: results from the Amsterdam Resuscitation Study (ARRESUST). *Resuscitation* 2001; 50:273-9.
 58. Deakin CD, O'Neill J F, Tabor T. Does compression-only cardiopulmonary resuscitation generate adequate passive ventilation during cardiac arrest? *Resuscitation* 2007; 75:53-9.
 59. Martens PR, Mullie A, Calle P, Van Hoeyweghen R. Influence on outcome after cardiac arrest of time elapsed between call for help and start of bystander basic CPR. The Belgian Cerebral Resuscitation Study Group. *Resuscitation* 1993; 25:227-34.
 60. Berg RA, Kern KB, Hilwig RW, et al. Assisted ventilation does not improve outcome in a porcine model of single-rescuer bystander cardiopulmonary resuscitation. *Circulation* 1997; 95:1635-41.
 61. Noc M, Weil MH, Tang W, Turner T, Fukui M. Mechanical ventilation may not be essential for initial cardiopulmonary resuscitation. *Chest* 1995; 108:821-7.
 62. Dorph E, Wik L, Stromme TA, Eriksen M, Steen PA. Oxygen delivery and return of spontaneous circulation with ventilation:compression ratio 2:30 versus chest compressions only CPR in pigs. *Resuscitation* 2004; 60:309-18.
 63. Dickinson ET, Verdile VP, Schneider RM, Salluzzo RF. Effectiveness of mechanical versus manual chest compressions in out-of-hospital cardiac arrest resuscitation: a pilot study. *Am J Emerg Med* 1998; 16:289-92.
 64. Wigginton JG, Miller AH, Benitez FL, Pepe PE. Mechanical devices for cardiopulmonary resuscitation. *Curr Opin Crit Care* 2005; 11:219-23.
 65. Ong ME, Ornato JP, Edwards DP, et al. Use of an automated, load-distributing band chest compression device for out-of-hospital cardiac arrest resuscitation. *JAMA* 2006; 295:2629-37.
 66. Bahr J. CPR education in the community. *Eur J Emerg Med* 1994; 1:190-2.
 67. Becker LB, Pepe PE. Ensuring the effectiveness of community-wide emergency cardiac care. *Ann Emerg Med* 1993; 22 (2 pt 2):354-65.